

Jet Studies with Higgs Events

Workshop on the Future of Higgs Physics

May 5, 2001

Sebastian Grinstein
for the DØ Collaboration

Tune jet algorithms parameters for Higgs searches:

- Introduction
- Calorimeter Tower Thresholds
- b -Jets Reconstruction Efficiencies
- b -Jets Energy Resolution
- Higgs Mass Resolution
- Conclusions

Introduction



- Preliminary studies of jet algorithms (Cone, kT) and algorithm parameters (R, D) effects on resolution and efficiency reconstruction in Higgs events.
- Full GEANT simulation of Run II detector.
[D0gstar (detector response) → D0Sim (digitalization) → D0reco (reconstruction)].
- Previous studies based on Run I detector simulation (hep-ph/0010338) or “fast” MC.
- RunII Jet Algorithms (kT, Cone) see hep-ex/0005012
- **Work in progress!**

Calorimeter Tower Threshold

Energy associated with average # of min bias events for that luminosity is subtracted from the event \Rightarrow contribution from min bias events: **Zero average \pm fluctuations**

Does the threshold of the calorimeter towers affect b jet resolution?

MC: $t\bar{t}$ with average 4.0 min bias overlaid (lowPt QCD)
 $E_{cal}/E_{particle}$ distributions \rightarrow




\rightarrow No Threshold





Selected Threshold: 0 MeV

Jets in Higgs Samples

MC Sample: $WH (W \rightarrow e\nu)$ 2.5mb ($10^{32}\text{cm}^{-2}\text{s}^{-1}$) [$m_H = 120 \text{ GeV}$]



Min $E_T = 8 \text{ GeV}$




For these events expect 3 jets ($b\bar{b}$ and e).

“Min Bias” Jets.


Jets in Higgs Samples

Same Event shown for Cone:


Parton




R=0.1



R=0.7



R=1.0



$R = 1.0$ “lost” a jet (by merging)

- We are in the process of understanding the Run II jet algorithms and studying their performance for Higgs searches.

b -Jets “Reconstruction” Efficiencies


WH ($W \rightarrow e\nu$) 2.5mb

b -quarks are matched in $\eta - \phi$ space to the reconstructed calorimeter jets. The “reconstruction” efficiency defined:

$$(Num_matched_calor_jets)/(Total_Num_of_b_quarks)$$

DR distributions ($\Delta R = \sqrt{\delta\eta^2 + \delta\phi^2}$)


between b -quarks and Calorimeter Jets (Cone):




Similar distributions for kT.
 $\Delta R \sim 0.2$ “reasonable” choice.

b-Jets “Reconstruction” Efficiencies

b Jet Reco Eff vs PT




b Jet Reco Eff vs PT



b -Jets “Reconstruction” Efficiencies


Cone Jets



- $R \sim 0.4$ provides the best overall “reconstruction” efficiency regardless the ΔR cut.
- Higher cone sizes sensitive to noise and smaller sizes affected by hadronization fluctuations.


b -Jets “Reconstruction” Efficiencies

kT Jets



- Similar behavior than Cone. $D \sim 0.3$ provides best overall “reconstruction” efficiency.

b-Jets Energy Resolutions




- WH ($W \rightarrow e\nu$) 2.5mb
- Matched Particle jets with calorimeter jets and calculate the variance of the $E_{cal}/E_{particle}$ distributions.
- Larger algorithm parameter choices yield poorer resolutions.

b-Jets Energy Resolutions

How much does the luminosity affect the resolutions?


- MC $t\bar{t}$ sample
- Resolution as a function of min bias overlaid.
- E_T range of 40 – 70 GeV



- Larger algorithm parameter sizes are more affected by luminosity.

Higgs Mass Resolution

- MC Sample: ZH ($Z \rightarrow \mu\mu$) 2.5mb, with initial and final state radiation [$m_H = 120$ GeV].
- b -quarks required to match calorimeter jets to simulate b tagging.
- Uncorrected calorimeter energies. Using reconstructed primary vertices.
- $M^2 = (E_1 + E_2)^2 - (p_{x1} + p_{x2})^2 - (p_{y1} + p_{y2})^2 - (p_{z1} + p_{z2})^2$




Preliminary results similar to previous studies with Run I detector simulation (hep-ph/0010338):

This study, Cone $R = 0.7$: $\sigma/M = 19.2\%$

Workshop 2000 [$m_H = 90$ GeV]: $\sigma/M = 17.6\%$

Higgs Mass Resolution

- MC Sample: WH ($W \rightarrow e\nu$) 2.5mb
($m_H = 120$ GeV).




- Better mass resolution for $R \sim 0.8$.


Higgs Mass Resolution

MC Sample: $ZH (Z \rightarrow \mu\mu)$
 + 2.5mb

If muon close to the calorimeter jet axis is “added in”
 \Rightarrow improves resolution.



Muon added to calorimeter jet.



Conclusions

- Preliminary results suggest Cone jets of $R \sim 0.5$ have good efficiency and resolution. Also used in Run I for top analyses.
- kT jets of $D \sim 0.4$ good efficiency and resolution compared to other D parameters.
- Higgs mass resolution results similar to those obtained for the “Higgs Workshop 2000” (hep-ph/0010338).
- All results are preliminary. Energy corrections and tracking information to be implemented soon.