

Beauty and Charm Physics at CDF Run II


Mary Bishai

(for the CDF II collaboration)

IV International Symposium on LHC Physics and
Detectors FERMILAB, May 1-3 , 2003

Topics



- b and c production - Run II
- Masses, lifetimes, branching fractions of b/c hadrons.
- CKM matrix physics at CDF





- New central tracker w/ 96 layers
- 8 Layers of 3-D Silicon up to $|\eta| = 2$, $\sigma(d_0) \sim 20\mu\text{m}$
- New TOF system
- Track Triggers at Level 1 (Central tracking) and Level 2 (Si tracking) with $p_t > 1.5 \text{ GeV}/c$ $|\eta| < 1.0$



Run II Secondary Vertex Trigger





$$150 \mu\text{m} \leq d_0 \leq 1 \text{mm} \quad 2^\circ \leq \Delta\phi \leq 90^\circ$$







Charm in Run II



Feb'-02 charm signals reconstructed with online SVT tracks.




450K $D^0 \rightarrow K\pi$ fully reconstructed 78K D^ fully reconstructed*



Direct Charm Production Run II





Use impact parameter of reconstructed charm mesons $F_D(d_0)$ to distinguish directly produced charm from $B \rightarrow DX$, $F_B(d_0)$







From $K_s \rightarrow \pi\pi$ data we find $F_D(d_0) = \text{Gaussian} + \text{exp tails}$.

From $B \rightarrow DX$ MC : $F_B(d_0) = \text{a double exponential}$.





Direct Charm Production X-Sec.




$E/T = 1.7 - 1.9$ (low p_t) and $E/T = 1.3 - 1.4$ (high p_t)



Theory update: B production



When $p_T \gg m_b$, large logarithms of the ratio p_T/m_b arise in the coefficients of the perturbative expansion. Resummation of next-to-leading logs merged with the QCD NLO using a retuned fragmentation function from more recent e^+e^- data with NLL applied



Cacciari, Nason hep-ph/0204025 (Run I Data)

$$E/T = 1.7 \pm 0.5(\text{theory}) \pm 0.5(\text{expt})$$



Run II: test current theory predictions at $p_t^{\text{min}} < 5 \text{ GeV}$



Run II Muon Triggers



Run II dimuon trigger has been optimized and running stably at a rate of $12 J/\psi s / nb^{-1}$ ($|\eta| < 1.0$) since Jan. 2002




L1 μ trigger eff. ($|\eta| < 0.6$, $plat.=98\%$) $p_t(J/\psi)$ GeV/c

Run II di-muon trigger features:


- lower p_t reach: $p_t(\mu) > 1.5(|\eta| < 0.6)$, $2.0(0.6 < |\eta| < 1.0)$
→ $p_t(J\psi) = 0$ GeV/c → TOTAL cross-sections
- Opening angle $\Delta\phi > 5^\circ$ (was 15°) → polarization.




Run II inclusive $J/\psi X$ x-sec



300K J/ψ from 37 pb^{-1}



Detector acceptance



J/ψ inclusive x-sec

First at a hadron collider !: Total J/ψ cross-section:

$$\begin{aligned} &\sigma(p\bar{p} \rightarrow J/\psi X, |y(J/\psi)| < 0.6) \\ &\times Br(J/\psi \rightarrow \mu\mu) \\ &= 240 \pm 1(stat)_{-28}^{+35}(syst) \text{ nb} \end{aligned}$$



HVQ production - summary



Run II new and improved triggers ! Larger kinematic coverage of the dimuon triggers, $|\eta| < 1$, $p_t(\mu) < 1.5 \text{ GeV}/c$ ($|\eta| < 0.6$). Track triggers \Rightarrow large SVT b/c yields at $p_t(l) < 6 \text{ GeV}/c$ and $p_t(\text{track}) > 2 \text{ GeV}/c$. *Test theory predictions in a different kinematic region.*

Milestones on the road to HVQ production: Preliminary measurements of total inclusive J/ψ and open charm x-secs. Exclusive B mesons reconstructed. methods for separation of prompt J/ψ and D from bs developed. *Preliminary measurements of b production x-secs underway.*


*Masses, Lifetimes, Branching
fractions*



SVXII Material calibration



J/ψ mass $\Rightarrow \sim 10\%$ of SVXII material unaccounted for. Need B-Field correction of $\approx 0.1\%$.







Particle	Measured mass	PDG 2002 mass
K_s	$497.36 \pm 0.05(stat)$	497.672 ± 0.031
D^0	$1864.25 \pm 0.06(stat)$	1864.6 ± 0.5
$\Upsilon(1S)$	$9458 \pm 1(stat)$	9460.37 ± 0.21



B meson masses



Run I B_s and Λ_b worlds best. Competitive measurements early.



Particle	Measured mass (GeV/c^2)	PDG 2002 mass
B^+	$5279.32 \pm 0.68(\text{stat}) \pm 0.94(\text{syst})$	5279.0 ± 0.5
B^0	$5280.30 \pm 0.92(\text{stat}) \pm 0.96(\text{syst})$	5279.4 ± 0.5
B_s^0	$5365.50 \pm 1.29(\text{stat}) \pm 0.94(\text{syst})$	5369.6 ± 2.4




Mass Difference $D_s^+ - D^+$



Measurement of $\Delta M(D_s^+ - D^+)$ is input to the overall PDG fit for all the charmed mesons. Predicted by HQET and Lattice QCD. $D_s^\pm, D^\pm \rightarrow \phi\pi, \phi \rightarrow KK$

First CDF Run II Publication!!

Effect	Syst. [MeV/c ²]
fitting	0.14
event selection	0.11
momentum scale	0.10
Tracker effects	0.06
Calibration procedure	0.03
Total	0.21



$$m(D_s^\pm) - m(D^\pm) = 99.41 \pm 0.38(stat) \pm 0.21(syst) \text{ MeV}/c^2$$



$$m(D_s^\pm) - m(D^\pm) = 99.2 \pm 0.5 \text{ MeV}/c^2 \text{ -PDG2002}$$





B_s lifetime using $B_s \rightarrow J/\psi\phi$



The ratio $\frac{\tau(B_s)}{\tau(B_d)}$ tests Heavy Quark Expansion theory predictions.



$B_u^+ \rightarrow J/\psi K^+$ control sample. $\tau(B^+) = 1.57 \pm 0.07(stat) \pm 0.02(syst)$ ps





$\tau(B_s) = 1.26 \pm 0.20(stat) \pm 0.02(syst)$ ps

$\tau(B_s) = 1.34 \pm 0.23(stat) \pm 0.05(syst)$ ps - CDF Run I





Beauty Baryons



$\Lambda_b \rightarrow \Lambda_c \pi \approx 73$ events (measure Br)

$\Lambda_b \rightarrow \Lambda_c \ell X \approx 600$ events



$$\tau_{\Lambda_b}(\Lambda_c \mu X) = xxx \pm 0.13(\text{stat}) \pm 0.07(\text{syst})\text{ps}$$




$$\tau_{\Lambda_b}(\Lambda_c \ell X) = 1.32 \pm 0.15(\text{stat}) \pm 0.07(\text{syst})\text{ps} - \text{CDF Run I}$$



Cabbibo suppressed D decay



Decays of D^0 to CP eigenstates K^+K^- and $\pi^+\pi^- \Rightarrow D^0$ mixing and CP violation. Use the π_s from $D^{*\pm} \rightarrow D^0\pi_s^\pm$ to tag flavor.



$$\frac{\Gamma(D^0 \rightarrow K^+K^-)}{\Gamma(D^0 \rightarrow K\pi)} = 9.38 \pm 0.18(stat) \pm 0.10(syst)\% \text{ CDF Run II}$$

$$\Gamma(D^0 \rightarrow K^+K^-)/\Gamma(D^0 \rightarrow K\pi) = 9.93 \pm 0.14 \pm 0.14\% \text{ FOCUS2003}$$

$$\frac{\Gamma(D^0 \rightarrow \pi^+\pi^-)}{\Gamma(D^0 \rightarrow K\pi)} = 3.686 \pm 0.076(stat) \pm 0.036(syst)\% \text{ CDF Run II}$$

$$\Gamma(D^0 \rightarrow \pi^+\pi^-)/\Gamma(D^0 \rightarrow K\pi) = 3.53 \pm 0.12 \pm 0.06\% \text{ FOCUS2003}$$


$$A_{cp}(D^0 \rightarrow KK) = 2.0 \pm 1.7(stat) \pm 0.6(syst)\%$$

$$A_{cp}(D^0 \rightarrow \pi\pi) = 3.0 \pm 1.9(stat) \pm 0.6(syst)\%$$




Rare FCNC $D^0 \rightarrow \mu\mu$


Sensitive to physics beyond the SM (3×10^{-13}). SUSY models with R-Parity violation predict up to 3.5×10^{-6}



Normalize to $D \rightarrow \pi\pi$



Pion misid rate



$$Br(D^0 \rightarrow \mu^+ \mu^-) \leq 2.4 \times 10^{-6} \quad @ 90\% \text{ C.L.}$$

$$\text{Previous limit} \leq 4.1 \times 10^{-6}$$




b/c -hadron properties



Particle	Measured mass MeV/c ²
B^+	$5280.6 \pm 1.7(stat) \pm 1.1(syst)$
B^0	$5279.8 \pm 1.9(stat) \pm 1.4(syst)$
B_s^0	$5360.3 \pm 3.8(stat)_{-2.9}^{+2.1}(syst)$
$m(D_s^\pm) - m(D^\pm)$	$99.41 \pm 0.38(stat) \pm 0.21(syst)$
Particle	Lifetime
B_s	$1.26 \pm 0.20(stat) \pm 0.02(syst)ps$
Λ_b	$xxx \pm 0.13(stat) \pm 0.07(syst)ps$
Decay	Branching fraction
$Br(D^0 \rightarrow \mu^+ \mu^-)$	$\leq 2.4 \times 10^{-6} @ 90\% C.L.$



Studying the α , β , γ of B Physics



B_s mixing ($\Delta m_s \gg \Delta m_d$)


Run II

Hadronic trigger

L00 and TOF

Use $B_s \rightarrow D_s^{(*)+} \underbrace{n\pi}_{\text{tag}}$


$$P_{SS(OS)}^{B_s^0} = 1/2\tau \exp(-t/\tau) \times [1 \mp \mathcal{A}(\Delta m_s) \cos(\Delta m_s t)]$$



Run I limit :

$$\Delta(m_s) > 5.8 \text{ps}^{-1} @ 95\% \text{ CL.}$$



From $B_s \rightarrow \phi l \underbrace{X\nu}_{\text{direction?}}$




$$X_s = \Delta m_s / \tau(B_s)$$



B_s hadronic - Run II



Compare $B_d \rightarrow D^- \pi$ and $B_u \rightarrow J/\psi K^+$ in TTT \Rightarrow trigger eff. understood. Normalize B_s to $B_d \rightarrow D^- \pi$, $D^- \rightarrow K \pi \pi$. B_s yield = 40 ± 10 events - much lower than expected but mostly understood.







Direct CPV in $b \rightarrow hh \rightarrow \alpha, \gamma$





Direct CP violation in $B^0 \rightarrow \pi^+\pi^-$ Penguin “pollution” need $B \rightarrow K^+K^-$. Expect 500 $B \rightarrow hh/100\text{pb}^{-1}$. Goal is to measure $A_{cp}(B_d \rightarrow \pi^+\pi^-, K^+\pi^- / B_s \rightarrow KK, K^-\pi^+ / \Lambda_b^0 \rightarrow p^+\pi^-, p^+K^-)$





Disentangling $B \rightarrow h^+ h^-$

Differences in $\pi - K$ kinematics separate $K\pi/\pi K$ from $\pi\pi/KK$ decays. Particle ID separates $\pi\pi$ from KK and πK from $K\pi$



$$M_{\pi\pi} \text{ versus } \alpha = \left(1 - \frac{p_1}{p_2}\right) \cdot q_1$$

$$id_{1,2} = \frac{dE/dX(trk_{1,2}) - dE/dX(\pi)}{dE/dX(K) - dE/dX(\pi)}$$

$$\mathcal{F} = G(M_{\pi\pi} - \mathcal{M}(\alpha), \sigma_M) \cdot \underbrace{P(\alpha)}_{MC} \cdot G(id_1 - E_{id_1}, 1) \cdot G(id_2 - E_{id_2}, 1)$$

$$\text{Total likelihood function : } \mathcal{L} = \sum_{i=1}^6 f_i \cdot \mathcal{F}_i(\alpha, M_{\pi\pi})$$



CONCLUSION



b/c Production *Key triggers and detector performances well understood. Preliminary measurements of the direct open-charm and total inclusive charmonium cross-sections (systematics limited).*

Masses, lifetimes, BRs *Worlds best B_s mass, lifetime and worlds largest samples of Λ_b with ONLY 65pb^{-1} . Current best limit on FCNC $D^0 \rightarrow \mu\mu$ decays*

CKM matrix physics: *Key hadronic signals $B_s \rightarrow D_s^{(*)}\pi$ and $B \rightarrow h^+h^-$ established. Initial rates with non-optimized SVT are lower than expected but now mostly understood.*



Backup slides





Detector performance



Status of L00





Clustering is well advanced. Now working on alignment and understanding resolution. Using L1 and L00 2 strip clusters to anchor track and look at residual in SVXII L0 - infer that without final alignment current 2-strip cluster resolution is $\sim 11\mu\text{m}$ (design is $8\mu\text{m}$).






Particle ID: TOF performance



Raw $\phi \rightarrow K^+ K^-$





$\phi \rightarrow K^+ K^- + \text{TOF}$



Lepton+SVT Track trigger Yields




By spring (100 pb^{-1}) CDF will have one of the worlds largest semileptonic B samples from the inclusive Lepton + SVT.



b decay	Expected yield
$b \rightarrow l + track$	3M
	(> 90% b-purity)
$B \rightarrow l D^0 X$	20K
	(B^\pm enhanced)
$B_s \rightarrow l D_s X$	1 K
$\Lambda_b \rightarrow \Lambda_c l \nu$	1 K



SVXII = Lots of Material



From comparison to GEANT MC, we find we are still missing $\sim 10\%$ of the SVXII material in the simulation.




Backup slides



HVQ Production Theory



Basics of HVQ Production in $p\bar{p}$



Two leading order perturbative processes:


$$\frac{\hat{\sigma}(gg \rightarrow Q_1\bar{Q}_1)}{\hat{\sigma}(gg \rightarrow Q_2\bar{Q}_2)} \xrightarrow{s \rightarrow \infty} 1 - \frac{\log(m_1^2/m_2^2)}{\log(s/m_2^2)}$$

$$\frac{\hat{\sigma}(q\bar{q} \rightarrow Q_1\bar{Q}_1)}{\hat{\sigma}(q\bar{q} \rightarrow Q_2\bar{Q}_2)} \xrightarrow{s \rightarrow \infty} 1 - \mathcal{O}(m_1^4/\hat{s}^2)$$

Hadronic states need PDFs:

$$\begin{aligned} \frac{d\sigma}{dyd\bar{y}dp_T^2} &\propto \frac{1}{m_T^4} \frac{1}{[1 + \cosh(y - \bar{y})]^2} \\ &\times \sum_{i,j} x_1 f_i(x_1) x_2 f_j(x_2) \\ &\times \sum_{\bar{}} |M(ij \rightarrow Q\bar{Q})|^2 \end{aligned}$$

Fragmentation \Rightarrow FS hadrons







Quarkonia production - Theory



Quarkonia bound states are *non-relativistic*. NRQCD LO perturbative expansion is $\mathcal{O}(\alpha_s^3 v^0)$ as in the color singlet model (CSM) + higher order $\mathcal{O}(\alpha_s^3 v^4)$. Fragmentation processes \propto color octet matrix element dominate.



Direct J/ψ production (Run I)

ψ' production (Run I)

At low p_t , non-fragmentation diagrams from other octet matrix elements are important, soft gluon effects cause rates to diverge.




Theory: B Fragmentation functions




Non-perturbative fragmentation functions for B mesons are extracted from LEP data using 3 different parameterizations.

Applied to LO and NLO QCD with \overline{MS} factorization. \Rightarrow good agreement with CDF Run I data on B meson cross-section.

Using the NRQCD factorization scheme: \Rightarrow good agreement with CDF Run I measurement of J/ψ cross-section from B .





Binnewies, Kniehl, Kramer hep-ph/9802231 (Run I)



hep-ph/9901348 (Run I)



b Production - Run II



Run II: $p_t(J/\psi) \sim 1.5\text{GeV} \Rightarrow p_t(B) \sim 0!$ Extract inclusive b-hadron cross-section from $b \rightarrow J/\psi X$ using long b lifetime



Λ_b and heavy baryon lifetimes



The dominant source of lifetime differences between b -flavored hadrons are weak interactions between the b -quark and the light valence quark of the order $16\pi^2(\bar{\Lambda}/m_b)^3 = \mathcal{O}(5 - 10\%)$ in HQET expansions where $\bar{\Lambda} = M_{H_b} - m_b$. The lifetime difference of b -hadrons therefore tests HQET expansions at the third order.

Expect $\frac{\tau(\Lambda_b)}{\tau(B_d)} \sim 0.9$ but measure $\frac{\tau(\Lambda_b)}{\tau(B_d)} = 0.78 \pm 0.05$.

Will new NLO QCD calculations explain the difference between theory and expt.?



Backup slides




CKM physics



Flavor Tagging



Tag B flavor at production:.





RUN II $\sin 2\beta$




RUN I

$$D_{\text{dilution}} = \frac{N_{\text{Right tags}} - N_{\text{Wrong tags}}}{N_{\text{Right tags}} + N_{\text{Wrong tags}}}$$


$$A_{\text{cp}}^{\text{observed}} = D A_{\text{cp}}^{\text{true}}$$

$$\sigma(\sin 2\beta) \sim 1 / \sqrt{\underbrace{\epsilon}_{\text{tag efficiency}} D^2 N_{\text{tot}}}$$



$$\sin(2\beta) = 0.91^{+0.37}_{-0.36}$$


Tag	RUN I ϵD^2	RUN II ϵD^2
JetQ	3.0%	3.0%
SLT	1.7%	1.7%
SST(π, K)	(1.4%, 1.0%)	(1.9%, 4.2%)
OS(K)	-	2.4%
Total (B_d)	6.1%	9.0%







Measuring $\sin \gamma$

$$\Gamma_{B_s^0 \rightarrow D_s^\mp K^\pm}(t) = Ae^{-t} [1 \pm R \cos x_s t \pm \sqrt{1 - R^2} \sin(x_s t) \sin(\delta + \gamma)]$$




$B_s^0 \rightarrow D_s^\mp K^\pm$ expected yield
and backgrounds:

$D_s \pi$ $D_s^* K$ $D_s^* \pi$



Expected error on $\sin(\delta + \gamma)$


B_d mixing and measuring $\sin 2\beta$



$$B(\bar{B}) \rightarrow l^\pm X, D^{*\mp} l^\pm \nu, l^\pm + jet$$

$$A(t) = \frac{\text{Unmixed}(t) - \text{Mixed}(t)}{\text{Unmixed}(t) + \text{Mixed}(t)} = \cos(\underbrace{\Delta m_d}_{M_{\text{heavy}} - M_{\text{light}}} t)$$

If $B\bar{B}$ mix AND decay to the same final state \Rightarrow CP



$$A_{CP} = \frac{N(\bar{B} \rightarrow f) - N(B \rightarrow f)}{N(\bar{B} \rightarrow f) + N(B \rightarrow f)} = \sin(2\beta) \sin(\Delta m_d t)$$



D^0 Mixing




(a) Lifetime difference between CP and non-CP final states

$$y \equiv \frac{\Delta\Gamma}{2\Gamma} = \frac{\tau_{K\pi}}{\tau_{KK,\pi\pi}} - 1$$

$$y < 10^{-5}$$

Expectation for 100 pb^{-1} . Belle/BaBar yield = $1M/100 \text{ fb}^{-1}$:

(b) Mixing in wrong sign hadronic decays = oscillations.



Mode	Yield
$D^0 \rightarrow K^- \pi^+$	1M
$D^0 \rightarrow K^+ K^-$	100K
$D^0 \rightarrow \pi^+ \pi^-$	30K
$D^0 \rightarrow K^+ \pi^-$ (DCS)	1 K



A_{cp} in $B_s \rightarrow J/\psi\phi$ and $B_s \rightarrow D_s^+ D_s^-$



Measures weak phase of V_{ts} which is very small in SM. Evidence for anomalous \mathcal{CP} phases if asymmetry is observed. $B_s \rightarrow D_s^- \pi^+$ is $\sim (50 : 50)$ CP even and odd, $B_s \rightarrow J/\psi\phi$ is mixed, $B_s \rightarrow D_s^+ D_s^-$ is CP even.

