

# W,Z + jets studies at Tevatron relevant to Higgs searches at LHC

R.Mazini (University of Toronto)

B.Mellado, W.Quayle and Sau Lan Wu  
(University of Wisconsin)



TEV4LHC, Higgs Session 09/16/04

---

# Outline

## + Introduction

- H+jets in Higgs searches at the LHC
- Considerations on background extraction


## + W,Z+jet studies for H+2jets (VBF)

## + Z+1jet studies for $\tau\tau$ +jet final state

## + Jet veto studies for $pp \rightarrow WW+X$

## + Outlook

# Low Mass SM Higgs Potential at LHC



# H+jets in Low Mass Higgs Searches at the LHC

H+jets will play a very important role at the LHC in observing a low mass Higgs

➤ These analyses are harder than purely inclusive ones

## Relative Sensitivity of H+jets (preliminary)

$H \rightarrow \tau\tau$	$H \rightarrow \gamma\gamma$	$H \rightarrow WW^{(*)}$	$H \rightarrow ZZ^{(*)}$
100%	~50%	~50%	~0% (?)


Together with  $t\bar{t}H(\rightarrow b\bar{b})$  most important final states close to LEP limit (114 GeV)

# On background extraction

- ✚ The determination of the SM (mostly QCD) backgrounds associated with jets will rely on LHC data itself
  - Will not rely on a prediction based on extrapolation from Tevatron to LHC
- ✚ Control samples in Data are well defined
  - Will require a certain degree of extrapolation from control sample phase space to signal-like region
  - If MC are used for this, extrapolation needs to be validated with Data
  - Tevatron plays a central role in validating MC tools, which will be extensively used at the LHC


# On background extraction (cont)

- Take example of  $H(\rightarrow\tau\tau)+\text{Jets}$ : Main background  $Z+\text{jets}$ 
  - Can be generalized to rest of final states



# H+2jets (VBF) at the LHC

✚ D.Zeppenfeld, D.Rainwater, et al. proposed to search for a Low Mass Higgs in association with two jets




# H+2jets (VBF) at the LHC (cont)

✚ Study additional (central) jet production to  $W + 2$  forward and separated jets (tagging jets)

- ❖ Cross-section dependence on separation in pseudorapidity between tagging jets
- ❖ Rate of third jet
- ❖ Angular correlations between tagging jets and central jet

➤ Comparison with QCD predictions

- ❖ Test interplay between perturbative and parton shower approaches





# H+2jets (VBF) at the LHC (cont)

## Effective cross-sections (in pb) evaluated with MadGraphII for the Tevatron

- Fast simulation with basic detector response (thanks to M.Martinez and Y-K.Kim)

Process	$\eta_{j_1 j_2} > 2$	$\eta_{j_1 j_2} > 2.5$	$\eta_{j_1 j_2} > 3$	$\eta_{j_1 j_2} > 3.5$	$\eta_{j_1 j_2} > 4$
QCD $Wjj$	7.07	4.29	2.40	1.23	0.56
EW $Wjj$	0.12	0.07	0.04	0.03	0.01
QCD $Wjjjj$	0.34	0.22	0.13	0.07	0.04

## W+2j and W+4j display large enough cross-section at the Tevatron

- Very hard to disentangle EW from QCD W/Z production


# H( $\rightarrow\tau\tau$ )+1jet at the LHC

✦ Tag one semi-central jet, require  $P_{TH} > 100$  and  $M_{HJ} > 700$  GeV and a loose central jet veto ("top killer")


➤ Allow significant contribution from  $gg \rightarrow h$

B. Mellado, W. Quayle  
and Sau Lan Wu  
hep-ph/0406095  
submitted to PL


Higgs Decay Products  
 $P_{TH} > 100$  GeV




- ✚ QCD  $Z+1j$  production gives about 50% of background
  - Need to evaluate role of QCD higher order corrections
    - ❖ These are not trivial due to specifics of cuts



# QCD HO Corrections in QCD Z+1jet



# K Factor





# QCD HO Corrections in QCD Z+1jet (Kinematic Effects)

- ✚ MC@NLO: Kinematic effects of Higher Order corrections to Z+1j (incorporates LO Z+1j ME) are given by Parton Showers
  - Certain limitations, which are analysis dependent
- ✚ SHERPA incorporates consistent matching between multi-parton tree-level ME with Parton showers
  - It gives rates to LO but it is a good tool to address kinematic effects of extra hard gluon radiation in the final state
  - Generate same number of events with Z+1j ME + PS and Z+2j ME + PS with SHERPA



# Z+1jet Study with SHERPA

Z+1j ME





Parton Shower for extra gluon radiation

Z+ $\leq 2$ j ME





Parton Shower for extra gluon radiation

- Require at least one jet with  $P_T > 30$  GeV
- To define an extra jet  $P_T > 30$  GeV is also required
  - $N_{cj}$  = number of extra jets with  $|\eta| < 2$







Strong effect on  $P_T$  of leading jet and the invariant mass of Z and the leading jet




Central jet veto ("top killer",  $P_{TJ} < 30\text{GeV}$ ) significantly reduces effect of higher order corrections



➤ With  $M_{ZJ} > 700\text{ GeV}$   $Z+1j$  increases by factor of 2



- ✦ Enhancement of  $Z+1j$  background due to extra gluon radiation may be excellent news to the analysis
- ✦  $H+1j$  and  $Z+1j$  diagrams bear strong similarities. Expect similar effect on Higgs production.
  - If signal enhancement turns out to be large then it would be a good idea to remove the central jet veto to further improve the signal significance
  - Requires study within SHERPA





- Require at least one jet with  $P_T > 20$  GeV
- To define an extra jet  $P_T > 20$  GeV is also required
  - $N_{cj}$  = number of extra jets with  $|\eta| < 2$




✚ Cross-section is large enough at the Tevatron to do studies at large  $P_{TZ}$ ,  $P_{TJ}$  and  $M_{ZJ}$

➤ About 100-200 fb for  $P_{TZ} > 100$  GeV




# W/Z + jets in CDF

Preliminary, (kindly provided by CDF)



**W+jets**



**Z+jets**  
(central electrons)


n	≥ 0	≥ 1	≥ 2	≥ 3	≥ 4
$N_n$	54799	11615	2680	602	145
$B_n$	1869	951	349	138	55

	≥ 0	≥ 1	≥ 2	≥ 3	≥ 4
$N_n^{cand}$	4232	935	193	40	4
$N_n^{bkg}$	6,51	8,01	3,82	0,67	0,10
$\epsilon_n(\%)$	80,53	76,66	70,31	69,78	85,76
$A_n(\%)$	10,81	10,13	9,40	9,48	11,70

# Jet veto studies for $pp \rightarrow WW + X$


✚ The application of a jet veto in  $pp \rightarrow WW + X$  is fundamental to Low Mass Higgs searches with  $H \rightarrow WW \rightarrow ll\nu\nu$  at the LHC

➤ MC will be validated with  $t\bar{t}, WW$  samples with  $M_{ll} > M_H$ , and ZW



✚ Different MC's (Pythia and MC@NLO/Herwig) predict very different  $P_T$  of  $t\bar{t}$  for LHC. Differences should be visible already at the Tevatron

➤  $P_T$  of  $t\bar{t}$  strongly correlated to jet multiplicity





# Outlook

- ✚ Higgs associated with jets play a central role in searches for Low Mass Higgs at the LHC
  - Need to extract reliably QCD backgrounds
    - ❖ Will rely on LHC data to extract QCD backgrounds
  - Tevatron plays a central role in validating MC tools, which will be extensively used at the LHC
- ✚ W/Z associated with jets are produced copiously enough at the Tevatron to study topologies relevant to H+1j and H+2j searches at the LHC
  - Cross-sections for W/Z+1,2,4 jets are large enough to investigate relevant corners of the phase-space
- ✚ Jet veto in  $pp \rightarrow WW+X$  is central to Higgs searches with  $H \rightarrow WW \rightarrow ll\nu\nu$  at the LHC